

# NONLINEAR RESPONSE OF SEMICONDUCTOR BRAGG REFLECTORS

V. V. Stankevich, M. V. Ermolenko, S. A. Tikhomirov, O. V. Bugarov,  
and S. V. Gaponenko

Institute of Molecular and Atomic Physics, NAS, Minsk 220072, Belarus.

A nonlinear periodic structure is proposed and developed with the purposeful use of strong resonant intrinsic interband absorption in a semiconductor material, which causes modification of the material refractive index in the spectral range corresponding to morphological resonances formed by the periodicity of the structure. In the experiments, periodic ZnSe/ZnS heterostructures have been used and interband excitation of a ZnSe sublattice has been performed by nano-, pico- and femtosecond laser pulses. A considerable shift of reflection spectrum and large relative reflection changes were observed in a wide spectral range corresponding to the transparency region of ZnSe far from the intrinsic absorption onset.

Evaluated refraction index change is about  $-0.05$ , the relaxation time being as short as 3 picoseconds. Relative reflection coefficient change exceeds 100%. In case of femtosecond excitation, a wide-band nonlinear response is observed for both one-photon near UV- and two-photon near IR-pulses. The nonlinearity relaxation time is supposed to depend a transition from non-equilibrium to quasi-equilibrium distribution of electrons and holes within ZnSe conduction and valence band, respectively, rather than electron-hole recombination time.

